

Contents Volume 81, 1994

<i>Preface</i>	i
Palaeobotanical investigation of early Tertiary palaeoelevations in northeastern Nevada: initial results D.A.R. Povey, R.A. Spicer and P.C. England	1
Atmospheric CO changes since the last glacial maximum: evidence from the stomatal density record of fossil leaves D.J. Beerling and W.G. Chaloner	11
Distinguishing internal developmental characteristics from external palaeoenvironmental effects in fossil wood J.L. Chapman	19
"Twig" - wood anatomical characters as palaeoecological indicators I. Poole	33
C enriched Lower Carboniferous fossil plants from Donegal, Ireland: carbon isotope constraints on taphonomy, diagenesis and palaeoenvironment T.P. Jones	53
A biomechanical method to distinguish between self-supporting and non self-supporting fossil plants T. Speck	65
The paleoecology of high-latitude Eocene swamp forests from Axel Heiberg Island, Canadian High Arctic D.R. Greenwood and J.F. Basinger	83
Patterns in tropical leaf litter and implications for angiosperm paleobotany R.J. Burnham	99
Latest occurrence of the extinct genus <i>Cedrelospermum</i> (Ulmaceae) in North America: <i>Cedrelospermum manchesteri</i> from Mexico S. Magallón-Puebla and S.R.S. Cevallos-Ferriz	115
First occurrence of <i>Czekanowskia</i> (Gymnospermae, Czekanowskiales) in the United States S. Ash	129
New data on the morphology and anatomy of the Devonian zosterophyll <i>Serrulacaulis</i> Hueber and Banks from Venezuela C.M. Berry and D. Edwards	141
Monosulcate angiosperm pollen from the López de Bertodano Formation (upper Campanian-Maastrichtian-Danian) of Seymour Island, Antarctica R.A. Askin	151
Multivariate analysis of Australian Eocene dispersed cuticle floras: influence of age, geography and taphonomy on biozonation A.I. Rowett and A.D. Sparrow	165
A new species of <i>Discinites</i> from the Lower Permian of China G. Zhifeng and B.A. Thomas	185
Earliest pollen-dominated microfloras from the early Late Carboniferous of the Tian Shan Mountains, NW China: their significance for the origin of conifers and palaeophytogeography Y.-X. Zhou	193
Further studies on a late Tournaisian (Lower Carboniferous) flora from Loch Humphrey Burn, Scotland: spore taxonomy and ultrastructure A.R. Hemsley, G. Clayton and J. Galtier	213
The pollen morphology of the tribes Loteae and Coronilleae (Papilionoideae: Leguminosae). 2, <i>Lotus</i> L. and related genera M.J. Diez and I.K. Ferguson	233
Pollen ultrastructure of <i>Hymenocardia</i> Walllich ex Lindley and comparison with other Euphorbiaceae D. Lobreau-Callen and M. Suarez Cervera	257
Ultrastructural and chemical studies on Oligocene fossil wood from Bovey Tracey, Devon, UK T.P. Jones	279
Taxonomy and stratigraphical distribution of <i>Verrucosporites nitidus</i> Playford, 1964 and related species E. Turnau, V.I. Avchimovitch, T.V. Byvscheva, G. Clayton, K.T. Higgs and B. Owens	289
Palynological observations on the "Nubian Sandstone", southwestern Libya A.O. Tekbali	297
Late-Quaternary changes of plant biomass as recorded by pollen-stratigraphical data: a discussion of the problem at Valle di Castiglione, Italy D. Magri	313

Pollen fossile de Leguminosae—Mimosoideae dans l'Oligocène inférieur du bassin de l'Ebre (Espagne)—implications paleoclimatiques et paleogeographiques	327
C. Cavagnetto and P. Guinet	327
<i>Malpophyllum</i> , a new name for the fossil genus <i>Palmophyllum</i> Conwentz, 1886	337
K.P.N. Kumaran	337
<i>Origin of Land Plants</i> , by L.E. Graham—R. Wayne	339

